

ALGEBRAIC RELATIONS FOR RECURSIVE SEQUENCES

Luigi Cimmino (cimmino@na.infn.it)

Abstract

Through the following, we establish the conditions which allow us to express recursive sequences of real numbers, enumerated through the recurrence relation $a_{n+1} = Aa_n + Ba_{n-1}$, by means of algebraic equations in two variables of degree $n \in N$. We do this, as far as we know, like it has never been formalized before.

I'd like to precise that the work was develop without the support of any well-know results about difference equations. Recently I discover that the present work has been treated already, but in a different context, at least in the two books [Car] and [Ela]. Never the less, I found it is a good idea to publish it. It would be a simple and ready-to-use instrument to they aren't familiar with difference equations.

1 Introduction

The famous Fibonacci's formula $a_{n+1} = a_n + a_{n-1}$ may be view as the simplest case of the more general one $a_{n+1} = Aa_n + Ba_{n-1}$. This kind of relations are usually employed in different disciplines, like economy, pure and applied mathematics, physic, biology and so on. So we may be interested about the conditions to be fulfilled when one wants to express a relation like that, in the flavour of algebraic formulas.

In what follows, we prove the statements concerning this equivalence; then, applying our results, we show as we can compute the domino tiling of the graph $W_4 \times P_{n-1}$ and, moreover, we prove rigorously that the recurrence relation $a_{n+1} = 6a_n - a_{n-1}$, starting with terms $a_0 = 1$ and $a_1 = 5$, enumerates the sequence of all Pythagorean triples (x, y, z) with $y = x + 1$, discussed by Mills[1].

2 Algebraic equation of degree n equals n -th term of the recursive sequence

Consider the recursive sequence $\{a_n\}$, with $n \in N$, obtained through the relation $a_{n+1} = Aa_n + Ba_{n-1}$ starting with a_0, a_1 .

If A, B are real number such that $A^2 + 4B > 0$ and x_-, x_+ are the real solution of equation $x^2 = Ax + B$, with $x_- < x_+$, then it results $a_n = ax_+^n + bx_-^n, \forall n \in N_0$, where $a = \frac{a_1 - a_0 x_-}{\sqrt{A^2 + 4B}}$ and $b = -\frac{a_1 - a_0 x_+}{\sqrt{A^2 + 4B}}$.

Proof. Consider the linear system of unknowns a and b :

$$\begin{aligned} ax_+ + bx_- &= a_1 \\ a + b &= a_0 \end{aligned}$$

It admits only one solution if $x_+ - x_- = \sqrt{A^2 + 4B}$ differs from zero. Now by solving the system, we obtain the solution

$$a = \frac{a_1 - a_0 x_-}{\sqrt{A^2 + 4B}} \quad \text{and} \quad b = -\frac{a_1 - a_0 x_+}{\sqrt{A^2 + 4B}}.$$

Let $f : n \in N_0 \rightarrow ax_+^n + bx_-^n$, so we have to prove that $f(n) = a_n, \forall n \in N_0$. This equality holds true for $n = 0, 1$. By induction, we assume

$$\begin{aligned} ax_+^{n-1} + bx_-^{n-1} &= a_{n-1} \\ ax_+^{n-2} + bx_-^{n-2} &= a_{n-2} \end{aligned}$$

then, by multiplying B to each side of the second equation, we have

$$aBx_+^{n-2} + bBx_-^{n-2} - Ba_{n-2} = 0.$$

Being $x_\pm^2 = Ax_\pm + B$ and replacing $B = x_\pm^2 - Ax_\pm$, one writes the latter as

$$\begin{aligned} (ax_+^{n-2})x_+^2 - (ax_+^{n-2})Ax_+ + (bx_-^{n-2})x_-^2 - (bx_-^{n-2})Ax_- - Ba_{n-2} &= \\ (ax_+^n + bx_-^n) - A(ax_+^{n-1} + bx_-^{n-1}) - Ba_{n-2} &= 0. \end{aligned}$$

and substituting $ax_+^{n-1} + bx_-^{n-1} = a_{n-1}$, we have

$$ax_+^n + bx_-^n = Aa_{n-1} + Ba_{n-2} = a_n.$$

■

Conversely, let us consider the following:

Proposition 1 *Let a, b, x_-, x_+ be real number such that $x_- < x_+$. Then, by setting $a_n = ax_+^n + bx_-^n, \forall n \in N_0$, the set $S = \{a_n\}_{n \in N_0}$ is recursively enumerable with starting terms $a_0 = a + b$, $a_1 = ax_+ + bx_-$ through recurrence relation $a_n = Aa_{n-1} + Ba_{n-2}$ with $A = x_+ + x_-$ and $B = -x_+x_-$.*

Proof. By hypotheses $a_0 = a + b, a_1 = ax_+ + bx_-$ and for $n \geq 1$, one has

$$\begin{aligned} a_n &= ax_+^n + bx_-^n = \\ &= (x_+ + x_-)(ax_+^{n-1} + bx_-^{n-1}) - x_+x_-(ax_+^{n-2} + bx_-^{n-2}) = \\ &= (x_+ + x_-)a_{n-1} - x_+x_-a_{n-2} \end{aligned}$$

For all $n \in N_0$ we have $a_n = (x_+ + x_-) a_{n-1} - x_+ x_- a_{n-2} = Aa_{n-1} + Ba_{n-2}$.
 ■

We have proved the following theorem

Theorem 2 *Let $\{a_n\}_{n \in N_0}$ be a sequence of real number. The following statements are equivalent*

$$a_n = Aa_{n-1} + Ba_{n-2} \quad (1)$$

$$a_n = ax_+^n + bx_-^n \quad (2)$$

$\forall n \in N_0$ where x_+ and x_- are the real solution of equation $x^2 = Ax + B$.

3 A recursive sequence of natural numbers

Before we present a first application to the graph theory, we prove two corollaries which result useful to play with sequences of natural numbers.

Corollary 3 *Under the hypotheses of Proposition 1, if $x_+ x_- < (x_+ + x_-) - 1$, then the sequence $\{a_n\}_{n \in N_0}$ is strictly increasing.*

Proof.

$$\begin{aligned} a_{n+1} &= (x_+ + x_-) a_n - x_+ x_- a_{n-1} > \\ &> (x_+ + x_-) a_n - [(x_+ + x_-) - 1] a_{n-1} = \\ &= (x_+ + x_-) [a_n - a_{n-1}] + a_{n-1} \geq \\ &\geq a_n - a_{n-1} + a_{n-1} = a_n \end{aligned}$$

that is, $a_{n+1} > a_n$, $\forall n \in N_0$. ■

Corollary 4 *Under the hypotheses of Corollary 3, if $a + b$, $ax_+ + bx_-$, $x_+ + x_-$ all belong to N and $x_+ x_- \in Z$, then $\forall n \in N_0$, $a_n \in N$.*

Proof. Let $a_0 = a + b$ and $a_1 = ax_+ + bx_-$, both belonging to N by hypotheses. One has

$$\begin{aligned} a_2 &= ax_+^2 + bx_-^2 = (x_+ + x_-) (ax_+ + bx_-) - x_+ x_- (a + b) = \\ &= (x_+ + x_-) a_1 - x_+ x_- a_0 \in N \end{aligned}$$

So, if

$$\begin{aligned} a_{n-1} &= ax_+^{n-1} + bx_-^{n-1} \in N \\ a_{n-2} &= ax_+^{n-2} + bx_-^{n-2} \in N \end{aligned}$$

then

$$\begin{aligned}
a_n &= ax_+^n + bx_-^n = \\
&= (x_+ + x_-) (ax_+^{n-1} + bx_-^{n-1}) - x_+x_- (ax_+^{n-2} + bx_-^{n-2}) = \\
&= (x_+ + x_-) a_{n-1} - x_+x_- a_{n-2} \in N
\end{aligned}$$

For all $n \in N_0$ we have $a_n \in N$. ■

Faase[4] proved that the number of domino tilings of the graph $W_4 \times P_{n-1}$, for each $n \geq 2$, can be enumerated through recurrence relation, which is given by the product of the n -th Fibonacci number and the n -th Pell number, *i.e.* $f_n p_n = (f_{n-1} + f_{n-2})(2p_{n-1} + p_{n-2})$, as Sellers[3] shown.

Given that the Fibonacci and Pell numbers are respectively defined as the sequences of integers

$$\begin{aligned}
f_0 &= 1, f_1 = 1 \text{ and } f_n = f_{n-1} + f_{n-2} \\
p_0 &= 1, p_1 = 2 \text{ and } p_n = 2p_{n-1} + p_{n-2}
\end{aligned}$$

applying theorem 2, we have

$$\begin{aligned}
f_n &= \frac{1}{\sqrt{5}} \left[\left(\frac{1+\sqrt{5}}{2} \right)^{n+1} - \left(\frac{1-\sqrt{5}}{2} \right)^{n+1} \right] \\
p_n &= \frac{1}{2\sqrt{2}} \left[(1+\sqrt{2})^{n+1} - (1-\sqrt{2})^{n+1} \right]
\end{aligned}$$

so, by multiplying them, we compute, as a function of n , the number of domino tilings of these graphs, $\forall n \geq 2$.

4 A further example

Now, let us consider the recurrence relation $a_{n+1} = 6a_n - a_{n-1}$, proposed by Mills[1]. The terms of the sequence, starting with $a_0 = 1$, $a_1 = 5$, are the third component of a Pythagorean triple with $y = x + 1$.

By applying theorem 2, we have

$$a_n = \frac{1}{4} \left[(\sqrt{2} + 2) (3 + 2\sqrt{2})^n - (\sqrt{2} - 2) (3 - 2\sqrt{2})^n \right].$$

where, by mean of corollary 4, a_n is in N , $\forall n \in N_0$.

For all odd natural number n , holds true the relation $n^2 = \left(\frac{n^2+1}{2} \right)^2 - \left(\frac{n^2-1}{2} \right)^2 = \left(\frac{n^2-1}{2} + 1 \right)^2 - \left(\frac{n^2-1}{2} \right)^2$, with $\left(\frac{n^2-1}{2} \right) \in N$. Hence, the numbers belonging to our sequence, enumerated through $a_k = 6a_{k-1} - a_{k-2}$ and

starting with odd a_0 and a_1 , are such that $\left(\frac{a_k^2-1}{2}\right) \in N$. Let us prove that $\forall k \in N_0$, $\frac{a_k^2-1}{2} = m(m+1)$, for suitable $m \in N$.

Proof. Let be $\frac{a_k^2-1}{2} = \left(\frac{\sqrt{2a_k^2-1}-1}{2}\right) \left(\frac{\sqrt{2a_k^2-1}+1}{2}\right)$. Thus, we have $\frac{\sqrt{a_k^2-1}-1}{2} = m_k \in N$ iff $2a_k^2 - 1 = d_k^2$, $d_k \in N$.
It follows that

$$a_k = \frac{1}{4} \left[\left(\sqrt{2} + 2 \right) \left(3 + 2\sqrt{2} \right)^k - \left(\sqrt{2} - 2 \right) \left(3 - 2\sqrt{2} \right)^k \right]$$

then, we have

$$\begin{aligned} 2a_k^2 - 1 &= 2 \left[\frac{1}{4} \left[\left(\sqrt{2} + 2 \right) \left(3 + 2\sqrt{2} \right)^k - \left(\sqrt{2} - 2 \right) \left(3 - 2\sqrt{2} \right)^k \right] \right]^2 - 1 = \\ &= \left[\frac{1}{2\sqrt{2}} \left[\left(2 + \sqrt{2} \right) \left(3 + 2\sqrt{2} \right)^k - \left(2 - \sqrt{2} \right) \left(3 - 2\sqrt{2} \right)^k \right] \right]^2 = \\ &= \left[\frac{1}{4} \left[\left(2\sqrt{2} + 2 \right) \left(3 + 2\sqrt{2} \right)^k - \left(2\sqrt{2} - 2 \right) \left(3 - 2\sqrt{2} \right)^k \right] \right]^2 \end{aligned}$$

that is to say

$$d_k = \frac{1}{2} \left[\left(\sqrt{2} + 1 \right) \left(3 + 2\sqrt{2} \right)^k - \left(\sqrt{2} - 1 \right) \left(3 - 2\sqrt{2} \right)^k \right].$$

By corollary 4, $\forall k \in N_0$, $d_k \in N$ and it results a sequence of odd natural numbers for $d_k = 6d_{k-1} - d_{k-2}$. So it follows $\frac{\sqrt{a_k^2-1}-1}{2} = \frac{d_k-1}{2} = m_k \in N$, that is $a_k^2 = \left(\frac{a_k^2-1}{2} + 1\right)^2 - \left(\frac{a_k^2-1}{2}\right)^2 = [m_k(m_k+1) + 1]^2 - [m_k(m_k+1)]^2 = m_k^2 + (m_k+1)^2$. ■

If we consider the d_k 's formula, one has

$$m_k = \frac{d_k - 1}{2} = \frac{1}{4} \left[\left(\sqrt{2} + 1 \right) \left(3 + 2\sqrt{2} \right)^k - \left(\sqrt{2} - 1 \right) \left(3 - 2\sqrt{2} \right)^k - 2 \right]$$

hence, letting k vary in N_0 , we have that every such triples are enumerated by mean of the form $(m_k, m_k + 1, a_k)$, where the m_k 's recurrence relation is given by

$$\begin{aligned} m_k &= \frac{1}{2} (6d_{k-1} - d_{k-2} - 1) = \frac{1}{2} (6d_{k-1} - d_{k-2} - 6 + 1 + 4) = \\ &= \frac{1}{2} [6(d_{k-1} - 1) - (d_{k-2} - 1) + 4] = \\ &= 6 \frac{(d_{k-1} - 1)}{2} - \frac{(d_{k-2} - 1)}{2} + 2 = 6m_{k-1} - m_{k-2} + 2 \end{aligned}$$

i.e. the empirical formula given by Hatch[5].

We have to prove that $(m_k, m_k + 1, a_k)$ are all the triples with $y = x + 1$. Then, we consider generic triple $(m, m + 1, c)$; from number theory there are $r, s \in \mathbb{N}$ relatively prime with different parity such that $r > s$ and $c = r^2 + s^2$. Moreover if m is even then $m = 2rs$ and $m + 1 = r^2 - s^2 = 2rs + 1$, otherwise $m = r^2 - s^2$ and $m + 1 = 2rs = r^2 - s^2 + 1$. We obtain two Pell's equations

$$\begin{array}{lll} \text{m even} & r^2 - s^2 - 2rs = 1 & \Rightarrow (r - s)^2 - 2s^2 = 1 \\ \text{m odd} & r^2 - s^2 - 2rs = -1 & \Rightarrow (r + s)^2 - 2r^2 = 1. \end{array}$$

If one assigns an index to r and s in each one of this cases, then they become $(r_i - s_i)^2 - 2s_i^2 = 1$ and $(r_j + s_j)^2 - 2r_j^2 = 1$. This kind of equation admits an infinity of solution; moreover, a suitable couple (r_o, s_o) is either solution of the first or of the second equation, never solution of both one. Now, we choose for even m_k , let i vary in the set of even natural numbers and for odd m_k , let j vary in the sets of all odd natural numbers. We take care they are well-ordered in both one of subsequences, therefore enumerating m_k so that it alternates even and odd index, we have to solve the equation $r_k^2 - s_k^2 - 2r_k s_k = (-1)^k$, which is $(r_k - s_k)^2 - 2s_k^2 = (-1)^k$. Setting $p_k = r_k - s_k$ and $q_k = s_k$, we have the equation $p_k^2 - 2q_k^2 = (-1)^k$, whose trivial solution is $p_0 = 1, q_0 = 0$ and fundamental one $p_1 = 1, q_1 = 1$; from fundamental solution we give each other solutions, since $(p_1 + \sqrt{2}q_1)^k (p_1 - \sqrt{2}q_1)^k = (-1)^k$, resolving system of two equations

$$\begin{aligned} p_k + \sqrt{2}q_k &= (1 + \sqrt{2})^k \\ p_k - \sqrt{2}q_k &= (1 - \sqrt{2})^k \end{aligned}$$

where clearly it holds $p_k + \sqrt{2}q_k > p_k - \sqrt{2}q_k$.

Now, subtracting and adding latter equations side to side, one has respectively

$$\begin{aligned} q_k = s_k &= \frac{1}{4} \left[\sqrt{2} (1 + \sqrt{2})^k - \sqrt{2} (1 - \sqrt{2})^k \right] \\ p_k &= \frac{1}{2} \left[(1 + \sqrt{2})^k - (1 - \sqrt{2})^k \right] \Rightarrow \\ \Rightarrow r_k &= \frac{1}{4} \left[(\sqrt{2} + 2) (1 + \sqrt{2})^k - (\sqrt{2} - 2) (1 - \sqrt{2})^k \right]. \end{aligned}$$

so, stated at proposition 1, we have $r_k = 2r_{k-1} + r_{k-2}$ and

$$\begin{aligned} s_k &= \frac{1}{4} \left[\sqrt{2} (1 + \sqrt{2}) (1 + \sqrt{2})^{k-1} - \sqrt{2} (1 - \sqrt{2}) (1 - \sqrt{2})^{k-1} \right] = \\ &= \frac{1}{4} \left[(\sqrt{2} + 2) (1 + \sqrt{2})^{k-1} - (\sqrt{2} - 2) (1 - \sqrt{2})^{k-1} \right] = r_{k-1} \end{aligned}$$

then it results

$$\begin{aligned} r_k^2 + s_k^2 &= (2r_{k-1} + s_{k-1})^2 + r_{k-1}^2 = 6(r_{k-1}^2 + s_{k-1}^2) - (r_{k-2}^2 + s_{k-2}^2) = \\ &= 6c_{k-1} - c_{k-2} = c_k. \end{aligned}$$

The last result involves that the two sets $\{a_k\}$ and $\{c_k\}$, being such that $a_0 = c_0 = 1$, $a_1 = c_1 = 5$ and having same recursive relation, are term by term identical. All triples with first and second components which mutually differ by one is in the form $(2r_n s_n, r_n^2 - s_n^2, a_n)$. To be precise, one has

$$\begin{aligned} 2r_n s_n &= \frac{1}{4} \left[(\sqrt{2} + 1) (3 + 2\sqrt{2})^k - (\sqrt{2} - 1) (3 - 2\sqrt{2})^n - (-1)^n 2 \right] \\ r_n^2 - s_n^2 &= \frac{1}{4} \left[(\sqrt{2} + 1) (3 + 2\sqrt{2})^n - (\sqrt{2} - 1) (3 - 2\sqrt{2})^n + (-1)^n 2 \right] \\ a_n &= \frac{1}{4} \left[(\sqrt{2} + 2) (3 + 2\sqrt{2})^n - (\sqrt{2} - 2) (3 - 2\sqrt{2})^n \right]. \end{aligned}$$

5 Conclusion

Before concluding, we note that

$$\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = x_+$$

which gives us the behavior of such sequences when n tends to infinity.

We have shown that the relation $a_{n+1} = Aa_n + Ba_{n-1}$ can be express as algebraic relation, so that we can directly compute the n -th term of such a sequence, without compute the preceding terms of it. Moreover, the sequence of triples with $y = x + 1$ gives us an example who shows how we can work using results we obtained, in the fashion of a powerful tool, to play with number theory.

References

- [Car] G.F. Carrier, C.E. Pearson, *Ordinary Differential Equations*, SIAM, 1991.
- [Ela] S.N. Elaydi, *An Introduction to Difference Equations*, Springer, 1996.
- [1] J.T.S. Mills, *Another Family Tree for Pythagorean Triples*, Mathematical Gazette 80, pp. 545-548, (November) 1995.
- [2] J.V. Uspensky and M.A. Heaslet, *Elementary Number Theory*, McGraw-Hill Book Company, New York and London, 1939.
- [3] J.A. Sellers *Domino Tilings and Product of Fibonacci and Pell Numbers*, Journal of Integer Sequences 5, 2002.

- [4] F.J. Faase, *On the Number of Specific Spanning Subgraphs of the Graphs $G \times P_n$* , Ars Combinatoria 49, pp. 129-154, 1998.
- [5] G. Hatch *Pythagorean Triples and Triangular Square Numbers*, Mathematical Gazette 79, pp. 51-55, (March) 1995.